

HEMITHYROIDECTOMY PARTIAL THYROIDECTOMY

Wound Care: Your wound will be covered with tape, which should be left in place for about 2 weeks. The tape will be removed at your first postoperative visit. You may notice some dried blood under the tape but that is of no concern. You can wash and shower with the tape in place and even get it wet. Once removed, the tape may need to be replaced and you will then need to change it every 2 or 3 days for the next 2 to 3 months in order to get the best possible cosmetic result. The preferred tape is narrow, flesh coloured Micropore which can be obtained from your chemist.

Activities: You should generally restrict vigorous activities for 1 or 2 weeks after surgery. Activities, which involve turning the head suddenly, such as driving in heavy traffic, should be avoided although local driving is acceptable. Commonsense is the best way to avoid straining your neck.

Local symptoms: A variety of local symptoms are common for several weeks after surgery including tightness, choking, or having to force down food. Headaches and tiredness are also common as is weakness of the voice with prolonged use. Occasionally a symptom may last for months. Swelling of the neck around the wound is also common and may benefit from daily massage with Vitamin E cream. Numbness of the skin above the wound may be present and may last for many months.

Late complications: The only delayed complication of concern is wound infection. This may have occurred if the wound becomes very red, hot and more swollen. If that occurs you must seek attention from your local doctor straight away who will arrange for you to have antibiotics.

Follow-up: generally your follow-up will consist of

- A visit at 2 to 3 weeks to have the tape removed. You will need to ring the office to make an appointment for that visit
- Final surgical check up at 3 months

In addition, if you have been placed on calcium supplements, you may need to be seen by your local doctor on a weekly basis to have a blood test to check your calcium level, and to have your calcium supplements progressively reduced (see "Calcium supplements" below). You will also need to be reviewed by your local doctor or endocrinologist at around 6 weeks and have thyroid function tests to check the function of your remaining thyroid. Your local doctor or endocrinologist may need to see you more frequently for specific review if there are any problems.

Country patients: If you are unable to return to Sydney for follow-up, it may be possible for the tape to be changed and for your follow-up to be undertaken by your local doctor. *Wecontinued back page*

**If you have been discharged on calcium medication please take this form and give it to your doctor one week after discharge.*

Dear Doctor

Your patient has been discharged on calcium supplements following their thyroid surgery. They have been asked to see you on a weekly basis to have their serum calcium level checked and their medication reduced according to the protocol:

On discharge they will normally be on:

Caltrate 2 twice a day

If calcium is normal at one week reduce to:

Caltrate 1 twice a day

If calcium is normal the next week reduce to:

Caltrate 1 daily

If calcium is normal the next week:

Cease Caltrate

If you have any questions or concerns, please contact your patient's surgeon as listed overleaf.

Would need to confirm these arrangements before you leave hospital.

Calcium supplements: A temporary drop in calcium levels may sometimes occur even after hemithyroidectomy or partial thyroidectomy. The calcium level is checked in hospital and if it is normal, you will be sent home on calcium supplements (Caltrate). You will then need to see your local doctor every week after discharge to have a blood test and to have the calcium levels checked. If, at each visit, the calcium level in the blood is normal, then the dose needs to be reduced according to the attached protocol that needs to be given to your local doctor.

If you have any problems, following your thyroid surgery, you should speak to your local doctor, who will arrange to contact your surgeon.

Professor Stan Sidhu PhD FRACS

Phone: 02 9437 1731
Mobile: 0400 817 023
E-mail: stansidhu@nebsc.com.au
Website: www.drstansidhu.com

**Suite 202, Level 2, AMA House
69 Christie Street
St Leonards NSW 2065**

ENDOCRINE SURGICAL UNIT

Professor Stan Sidhu
www.drstansidhu.com

Postoperative Instructions

Royal North Shore Hospital
Mater Hospital
Sydney Adventist Hospital
Hornsby Hospital

HEMITHYROIDECTOMY PARTIAL THYROIDECTOMY



Suite 202/69 Christie St
AMA House
St Leonards NSW 2065